

# **Biological Alteration of Sedimentary Event-Layers on the Eel River Shelf: Phase 2**

Robert A. Wheatcroft  
College of Oceanic & Atmospheric Sciences  
Oregon State University  
Corvallis, OR 97331  
phone: (541) 737-3891 fax: (541) 737-2064 e-mail: [raw@oce.orst.edu](mailto:raw@oce.orst.edu)  
Award #: N00014-95-1-0165

## **LONG-TERM GOAL**

The ultimate objective of this research program is to obtain a predictive understanding of the physical and biological processes responsible for the formation, alteration and preservation of marine sedimentary deposits. While present focus is on the biological alteration of sedimentary event layers, the physical processes leading to the deposition and erosion of marine sediment remains an area of interest. The general approach of this research program is the development and testing of theory mainly through field observations and measurements.

## **OBJECTIVES**

The focus of this project, which is a component of the STRATAFORM program, is the documentation of the post-depositional biological alteration of sedimentary event layers (i.e., beds derived from wind-driven storms and river floods) on the Eel River shelf off northern California. Specifically, the spatial (along and across-isobath) and temporal variation of the small-scale geometry of event-bed contacts is quantified using digital x-radiography. Variations in geometry will be explicitly related to spatial and temporal patterns of independently measured forcings, such as bioturbation intensity, macrofaunal community structure, sediment accumulation rate and event bed thickness

## **APPROACH**

Box cores are the primary sampling device used in this research. Cores are taken in two different modes: (1) replicate time-series sampling of four stations along the 70-m isobath, and (2) broad, large-scale coverage of the Eel margin. Subsequent sources of data include transmission x-radiographs, microresistivity profiles, profiles of the naturally occurring radionuclides, Pb-210 and Th-234, and macrofaunal community composition, abundance and biomass. X-radiographs are digitized on a flatbed scanner and analyzed using a variety of image processing techniques (e.g., enhancement, segmentation).

## **WORK COMPLETED**

During FY 98, I was chief scientist on three STRATAFORM cruises (WR9710, PS9803 and leg 1 of W9807). During the first two cruises, replicate time-series samples were collected at four stations along the 70-m isobath representing areas severely to moderately impacted by the January 1995 and January 1997 flood deposits. The third cruise primarily focused on collecting (during transit) box and piston cores off other major river systems (e.g., Rogue, Klamath) in the Pacific Northwest.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>1998</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1998 to 00-00-1998</b>	
4. TITLE AND SUBTITLE <b>Biological Alteration of Sedimentary Event-Layers on the Eel River Shelf: Phase 2</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Oregon State University ,College of Oceanic and Atmospheric Sciences,Corvallis,OR,97331</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM002252.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>3</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## RESULTS

Analyses during FY 98 focused on the time-series data collected over the past three years. A key finding is that despite greater than 15 cm of sediment accumulation over a 3 year period at some locations, the shelf macrofauna display remarkable resiliency. That is, it is difficult to recognize a flood-effect following the 1995 and 1997 events, except for a short-term decline in the abundance of small, tube-dwelling polychaetes (e.g., spionids). Within-sediment depth distribution is similarly unaffected by the flood deposits (i.e., animals do not appear to avoid the deposits). Lastly, sediment mixing intensity is relatively high ( $\sim 50 \text{ cm}^2/\text{yr}$ ) and insensitive to the flood events. All of these data indicate that biological reworking on the mid-shelf is relatively constant in time.

A note-worthy feature of the Eel River shelf fauna, compared to that found on other West Coast shelves, is the dominance of polychaete worms and near complete absence of echinoderms (e.g., heart urchins and brittle stars) (Table 1). Presently, the explanation for this pattern is unknown; however, it may be a result of a local extirpation due to the 1964 flood (a 400-year return period event). That is, due to their extremely slow growth rates coupled with frequent floods over the past 35 years, the echinoderms have never recovered. Regardless of the cause, the absence of heart urchins has important ramifications for the preservation of event beds.

---

**Table 1. Percent fraction of major phyla (1-mm sieve fraction)**

Phylum	Site				
	WA	OR	Eel	Russian	LA
Annelida (worms)	48.2	55.1	86.1	61.8	65.6
Mollusca (clams, snails)	27.0	38.0	7.8	17.4	< 1
Arthropoda (crustaceans)	18.1	3.2	5.7	2.2	13.0
Echinodermata	6.6	3.6	< 1	18.5	21.6

Sources: WA (Lie and Kisker, 1970), OR (Richardson et al., 1977); Russian (C.A. Butman, WHOI, unpublished data); LA (Wheatcroft and Martin, 1996)

---

## IMPACT/APPLICATIONS

Documenting the initial distribution and subsequent modification of sedimentary event beds will provide key insight for modelers of strata development on continental margins.

## **TRANSITIONS**

Sedimentary strata have important and diverse acoustical implications. In a preliminary exploration of those implications, a selection of digitized x-radiographs have been subject to a statistical analysis in both the spatial and wave number domains by Dr. Dajun Tang (APL-University of Washington).

## **RELATED PROJECTS**

Field sampling has been a joint effort with Drs. J. Borgeld (HSU), D. Drake (USGS), C. Nittrouer (UW) and C. Sommerfield (WHOI). In addition, collaborations have been initiated with Dr. P. Wiberg (UVA) to examine the formation and preservation of the flood deposits.

## **REFERENCES**

Lie, U., and D.S. Kisker. 1970. Species composition and structure of benthic infauna communities off the coast of Washington. *J Fish. Res. Bd. Canada* 27: 2273-2285.

Richardson, M. D., A. G. Carey and W.A. Colgate. 1977. Aquatic disposal field investigations: Columbia River disposal site, Oregon. Tech. Rep. D-7730, USACE Waterways Experimental station, 411 pp.

Wheatcroft, R. A. and W. R. Martin. 1996. Spatial variation in short-term (Th-234) sediment bioturbation intensity along an organic carbon gradient. *J. Marine Res.*, 54: 763-792.